

DESCRIPTION

Fenbendazole is a broad spectrum benzimidazole anthelmintic used against gastrointestinal parasites including roundworms, hookworms, whipworms, the taenia species of tapeworms, pinworms, aerulostrongylus, paragonimiasis, strongyles and strongyloides.

In cattle and sheep, fenbendazole is active against *Dictiocaulus viviparous* and also against 4th stage larvae of *Ostertagia* spp. Fenbendazole also has an ovicide action. Fenbendazole acts by disrupting the the formation of microtubuli by binding to tubulin in the parasitic intestinal cells, preventing absorption of glucose. Fenbendazole is poorly absorbed after oral administration, reaching max after 20 hours in ruminants and more rapidly in monogastics. It is metabolized by the liver, and excreted within 48 hours in the faeces, and only 10% in the urine.

COMPOSITION

Fenbendazole 22.20 mg/g



INDICATIONS

Cattle: treatment of infestations by adult and immature forms of gastro-intestinal and respiratory nematodes. Also active against inhibited larvae of *Ostertagia* spp. and against *Moniezia* spp. of tapeworms.

Sheep: treatment of infestations by adult and immature forms of gastro-intestinal and respiratory nematodes. Also active against *Moniezia* spp. and useful but with variable efficacy against *Trichuris* spp.

Horses: for treatment and control of adult and immature stages of gastrointestinal roundworms in horses and other equidae.

Pigs: treatment of infestations by mature and immature nematodes of the gastro-intestinal tract, and control of roundworms in the respiratory tract and their eggs.

CONTRA-INDICATIONS

None.

DOSAGE AND MODE OF ADMINISTRATION

For oral administration.

Standard dose for ruminants and pigs is 5 mg fenbendazole per kg b.w. (=1 g HUNTER 22 per 40 kg b.w.).

For horses and other equidae, use 7.5 mg fenbendazole per kg b.w. (= 10 g HUNTER 22 per 300 kg b.w.).

Administer with the feed or on top of the feed.

SIDE EFFECTS

None.

SPECIAL PRECAUTIONS AND WARNINGS

Assess bodyweight as accurately as possible before calculating dosage. Direct contact with skin should be kept to a minimum. Wash hands after use.

PACK SIZE

100g, 200g, 500g, 1kg, 5kg



Manufacturer: Hebei Weierli Animal Pharmaceutical Group Co., Ltd

Exporter: Hebei Nuob Trade Co., Ltd

Website: www.victorypharm.com

Alibaba: www.victorypharm.en.alibaba.com

Email: julia@victorypharm.com

Mobie&Whatsapp: +86 13191871625

Tel: +86-311-89265710 **Fax:** +86-311-58052865



Address: B-1601, Lechen Business Building, No.260 Huaian Road, Shijiazhuang City, Hebei, China