

COMPOSITION

Each chewable tablet contains:

Praziquantel USP.... 50 mg

Pyrantel Pamoate USP... 144 mg

Febantel BP... 150 mg

Excipients...q.s.



INDICATION:

For the control of the following gastrointestinal tapeworms and roundworms of dogs and puppies.

Ascarids : *Toxocara Canis*, *Toxascaris leonine* (adult and late immature forms).

Hookworms : *Uncinaria stenocephala*, *Ancylostoma caninum* (adults).

Whipworms : *Trichuris vulpis* (adults).

Tapeworms: Echinococcus species, Taenia species, *Dipylidium caninum* (adults and immature forms).

Dosage: For oral administration only.

The recommended dose rates are: 15 mg/kg bodyweight febantel, 14.4 mg/kg pyrantel peanut and 5 mg/kg praziquantel. - 1 Febantel Plus Chewable Tablet per 10 kg bodyweight;

The tablets can be given directly to the dog or disguised in food. No starvation is needed before or after treatment. - For routine control adult dogs should be treated every 3 months. For routine treatment a single dose is recommended. In the event of heavy roundworm infestations repeat dose should be given after 14 days.

DOSAGE

Puppies and small dogs

1/2-2 kg bodyweight = 1/4 Tablet

2-5 kg bodyweight = 1/2 Tablet



5-10 kg bodyweight = 1 Tablet

Medium dogs

10-15 kg bodyweight = 1½ Tablet

15-20 kg bodyweight = 2 Tablet

20-25 kg bodyweight = 2½ Tablet

25-30 kg bodyweight = 3 Tablet

- For routine treatment a single dose is recommended.

Puppies should be treated at 2 weeks of age and every 2 weeks until 12 weeks of age. Thereafter they should be treated at 3 month intervals. It is advisable to treat the bitch at the same time as the puppies.

- For the control of Toxocara, nursing bitches should be dosed 2 weeks after giving birth and every two weeks until weaning.

Use During Pregnancy and Lactation:

- Consult a veterinary surgeon before treating pregnant animals for roundworms.

- The product may be used during lactation.

- Do not exceed recommended dose when treating pregnant bitches.

Contraindications, warnings, etc.:

- Do not use simultaneously with piperazine compounds.

- User safety: In the interests of good hygiene, persons administering the tablets directly to the dog, or by adding them to the dog's food, should wash their hands afterwards.

Storage: Store below 25°C.

PACKING

10 Tablets/ Blister, 10 Blisters/Box, 40 Boxes/Carton

100 Tablets/Satchel, 1 Satchel/Jar, 60 Jars/Carton

KEEP OUT OF REACH OF CHILDREN

For Veterinary Use Only.



Not for human use - for treatment of animals only.



Manufacturer: Hebei Pude Animal Medicine Co., Ltd

Exporter: Hebei Nuob Trade Co., Ltd

Website: www.victorypharm.com

Alibaba: www.victorypharm.en.alibaba.com

Email: julia@victorypharm.com

Mobie&Whatsapp: +86 13191871625

Tel: +86-311-89265710 **Fax:** +86-311-58052865



Address: B-1601, Lechen Business Building, No.260 Huaian Road, Shijiazhuang City, Hebei, China