

COMPOSITION

Each ml contains 4.54 mg of Pyrantel base as Pyrantel Pamoate.

INDICATION:

Treatment for the removal of large roundworms

and hookworms in dogs and puppies.

It is also effective in preventing the reinfestation of

T. canis in adult dogs, puppies and lactating moms after whelping.

DOSAGE AND ADMINISTRATION:

Administer 1 teaspoon (5 ml) for each 10 lbs of body weight. It is not necessary to withhold food prior to or after treatment. Dogs usually find this dewormer very palatable and will lick the dose from the bowl willingly. If there is reluctance to accept the dose, mix in a small quantity of dog food to encourage consumption. It is recommended that dogs under constant exposure to worm infestation should have a follow-up fecal exam within 2 to 4 weeks after treatment.

For maximum control and prevention of reinfestation, it is recommended that puppies be treated at 2, 3, 4, 6, 8, and 10 weeks of age. Lactating bitches should be treated 2 to 3 weeks after whelping. Adult dogs kept in heavily contaminated quarters may be treated monthly.

STORAGE:

Below 30°C (86°F).

PACKING:

30ml, 55ml, 100ml

CAUTION:

For use in dogs only.

Shake liquid thoroughly before every use.

The liquid is a suspension and will separate. To ensure uniform resuspension and to achieve







WORMER

Pyrantel Pamoate Suspension

indworn

proper dosage, it is important that the product be shaken thoroughly before every use.

If dog looks or acts sick, do not treat with this product



Manufacturer: Hebei Pude Animal Medicine Co., Ltd

Exporter: Hebei Nuob Trade Co., Ltd

Website: www.victorypharm.com

Alibaba: www.victorypharm.en.alibaba.com

Email: julia@victorypharm.com

Mob&Whatsapp&Wechat: +86 18332395825

Tel: +86-311-89265710 Fax: +86-311-58052865

Address: B-1601, Lechen Business Building, No.260 Huaian Road, Shijiazhuanng City, Hebei, China



